

**Anti-Murine B7-1 Antibody**  
**Catalog # ABG10016****Specification**

---

**Anti-Murine B7-1 Antibody - Product Information**

Application	<b>WB, E</b>
Reactivity	<b>Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

**Anti-Murine B7-1 Antibody - Additional Information****Preparation**

Produced from sera of rabbits immunized with highly pure recombinant Murine B7-1. Anti-Murine B7-1 specific antibody was purified by affinity chromatography employing an immobilized Murine B7-1 matrix.

**WesternBlot**

To detect mB7-1 by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mB7-1 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

**Sandwich**

To detect mB7-1 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Murine B7-1 (61-401BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mB7-1.

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Reconstitution**

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Murine B7-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-Murine B7-1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-Murine B7-1 Antibody - Images**