

Anti-Human CNTF Antibody

Catalog # ABG10047

Specification

Anti-Human CNTF Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human CNTF Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hCNTF. Anti-Human CNTF specific antibody was purified by affinity chromatography employing immobilized hCNTF matrix.

WesternBlot

To detect hCNTF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hCNTF is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hCNTF by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human CNTF (60-105BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hCNTF.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human breast invasive ductal carcinoma. The recommended concentration is 0.25 μ g/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Optimal results for these conditions were achieved without antigen retrieval. Optimal concentrations and conditions may vary.

Neutralization

To yield one-half maximal inhibition [ND_{>50}] of the biological activity of Human CNTF ($1.0 \mu g/ml$) a concentration of $7.0 - 11.0 \mu g/ml$ of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions



Anti-Human CNTF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human CNTF Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human CNTF Antibody - Images