

# **Anti-Rat CNTF Antibody**

Catalog # ABG10048

# **Specification**

# **Anti-Rat CNTF Antibody - Product Information**

Application WB, IHC, E
Reactivity Rat
Host Rabbit
Clonality Polyclonal

## **Anti-Rat CNTF Antibody - Additional Information**

#### **Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant Rat CNTF. Anti-Rat CNTF specific antibody was purified by affinity chromatography employing immobilized Rat CNTF matrix.

#### WesternBlot

To detect Rat CNTF by Western Blot analysis this antibody can be used at a concentration of  $0.1\text{-}0.2~\mu\text{g/ml}$ . Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat CNTF is 1.5-3.0~ng/lane, under either reducing or non-reducing conditions.

#### Sandwich

To detect Rat CNTF by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Rat CNTF (62-052BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat CNTF.

## **Immunohistochemistry**

This antibody stained colchicine injected rat brain (including the globus pallidus region) tissue. The primary antibody was incubated at 0.25 mg/ml overnight at  $4^{\circ}$ C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA $^{\text{TM}}$ ) reagent. Optimal concentrations and conditions may vary. <em><span style="font-size: 10px;"> <strong>Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm. </strong></span></em>

#### **Neutralization**

To yield one-half maximal inhibition [<strong>ND</strong><span style="font-size: 16px;"><sub>>50</sub></span>] of the biological activity of Rat CNTF (0.15 ng/ml), a concentration of 0.02-0.05 µg/ml of this antibody is required.

### **Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

#### Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

#### **Storage**

-20°C



Tel: 858.875.1900 Fax: 858.875.1999

# **Precautions**

Anti-Rat CNTF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# **Anti-Rat CNTF Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

**Anti-Rat CNTF Antibody - Images**