10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Human CXCL16 Antibody

Catalog # ABG10057

Specification

Anti-Human CXCL16 Antibody - Product Information

Application	**WB, IHC, E**
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**

Anti-Human CXCL16 Antibody - Additional Information

Preparation
Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hCXCL16. Anti-Human CXCL16 specific antibody was purified by affinity chromatography employing immobilized hCXCL16 matrix.

WesternBlot
To detect hCXCL16 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hCXCL16 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich
To detect hCXCL16 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human CXCL16 (60-109BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hCXCL16.

Immunohistochemistry
This antibody stained formalin-fixed paraffin-embedded sections of human colorectal adenocarcinoma tissue. The recommended concentration is 0.5 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Antigen retrieval is not required for these conditions. Optimal concentrations and conditions may vary.

 *Additional Immunostaining data available. Please contact Tech Support for information.

Formulation
A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution
Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage
-20°C

Precautions
Anti-Human CXCL16 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Human CXCL16 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human CXCL16 Antibody - Images