

Biotinylated Anti-Human EGF Antibody

Catalog # ABG10066

Specification

Biotinylated Anti-Human EGF Antibody - Product Information

Application WB, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Biotinylated Anti-Human EGF Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hEGF. Anti-Human EGF specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot

To detect hEGF by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hEGF is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hEGF by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.25 – 1.0 μ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti-Human EGF (60-115P) as a capture antibody, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hEGF.

Direct

To detect hEGF by direct ELISA (using 100 μ l/well antibody solution) a concentration of 0.25 – 1.0 μ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hEGF.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.


Storage

-20°C

Precautions

Biotinylated Anti-Human EGF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Biotinylated Anti-Human EGF Antibody - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Biotinylated Anti-Human EGF Antibody - Images