

Anti-Human ENA-78 Antibody
Catalog # ABG10076**Specification**

Anti-Human ENA-78 Antibody - Product Information

| | |
|-------------|-------------------|
| Application | WB, IHC, E |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |

Anti-Human ENA-78 Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hENA-78. Anti-Human ENA-78 specific antibody was purified by affinity chromatography employing immobilized hENA-78 matrix.

WesternBlot

To detect hENA-78 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hENA-78 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hENA-78 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human ENA-78 (60-118BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hENA-78.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human colon/rectum adenocarcinoma. The recommended concentration is 1.0 µg/ml-2.0 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human ENA-78 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human ENA-78 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human ENA-78 Antibody - Images