

Anti-Human FGF-4 Antibody

Catalog # ABG10107

Specification

Anti-Human FGF-4 Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human FGF-4 Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hFGF-4. Anti-Human FGF-4 specific antibody was purified by affinity chromatography employing immobilized hFGF-4 matrix.

WesternBlot

To detect hFGF-4 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hFGF-4 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hFGF-4 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human FGF-4 (60-134BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hFGF-4.

Immunohistochemistry

This antibody stained formalin-fixed paraffin-embedded sections of normal breast tissue from a breast carcinoma. The recommended concentration is 0.25 μ g/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 Sodium Citrate buffer is recommended. Optimal concentrations and conditions may vary.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human FGF-4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.





Anti-Human FGF-4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human FGF-4 Antibody - Images