10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Rat GRO-β/MIP-2 Antibody

Catalog # ABG10149

Specification

Anti-Rat GRO-β/MIP-2 Antibody - Product Information

Application	**WB, E**
Reactivity	**Rat**
Host	**Rabbit**
Clonality	**Polyclonal**

Anti-Rat GRO-β/MIP-2 Antibody - Additional Information

Preparation
Produced from sera of rabbits pre-immunized with highly pure recombinant Rat GROβ/MIP-2. Anti-Rat GROβ/MIP-2 specific antibody was purified by affinity chromatography employing immobilized Rat GROβ/MIP-2 matrix.

WesternBlot
To detect Rat GROβ/MIP-2 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Rat GROβ/MIP-2 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich
To detect Rat GROβ/MIP-2 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Rat GROβ/MIP-2 (62-060BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat GROβ/MIP-2.

Neutralization
To yield one-half maximal inhibition [ND₅₀] of the biological activity of Rat GROβ/MIP-2 (100 ng/ml), a concentration of 3.0-5.0 µg/ml of this antibody is required.

Formulation
A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution
Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage
-20°C

Precautions
Anti-Rat GRO-β/MIP-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Rat GRO-β/MIP-2 Antibody - Protocols

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Rat GRO-β/MIP-2 Antibody - Images