

Anti-Human IL-17F Antibody

Catalog # ABG10229

Specification

Anti-Human IL-17F Antibody - Product Information

Application WB, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human IL-17F Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-17F. Anti-Human IL-17F specific antibody was purified by affinity chromatography employing immobilized hII-17F matrix.

WesternBlot

To detect hIL-17F by Western Blot analysis this antibody can be used at a concentration of 0.1 - $0.2 \mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-17F is 1.5 - $3.0 \mu g/lane$, under either reducing or non-reducing conditions.

Sandwich

To detect hIL-17F by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human IL-17F (60-017FBT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hIL-17F.

Neutralization

To yield one-half maximal inhibition [ND₅₀] of the biological activity of hIL-17F (50.0 ng/ml), a concentration of 0.9-1.2 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.


Storage

-20°C

Precautions

Anti-Human IL-17F Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human IL-17F Antibody - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IL-17F Antibody - Images