

Biotinylated Anti-Human IL-1α Antibody

Catalog # ABG10238

Specification

Biotinylated Anti-Human IL-1α Antibody - Product Information

Application WB, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Biotinylated Anti-Human IL-1α Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hIL-1 α . Anti-Human IL-1 α specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot

To detect hIL- 1α by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL- 1α is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hIL- 1α by sandwich ELISA (using $100 \, \mu$ l/well antibody solution) a concentration of $0.25 - 1.0 \, \mu$ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti-Human IL- 1α (60-001AP) as a capture antibody, allows the detection of at least $0.2 - 0.4 \, n$ g/well of recombinant hIL- 1α .

Direct

To detect hIL-1 α by direct ELISA (using 100 μ l/well antibody solution) a concentration of 0.25 – 1.0 μ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hIL-1 α .

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage


-20°C

Precautions

Biotinylated Anti-Human IL- 1α Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Biotinylated Anti-Human IL-1α Antibody - Protocols







Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Biotinylated Anti-Human IL-1α Antibody - Images