

Biotinylated Anti-Murine IL-22 Antibody

Catalog # ABG10267

Specification

Biotinylated Anti-Murine IL-22 Antibody - Product Information

Application WB, E
Reactivity Mouse
Host Rabbit
Clonality Polyclonal

Biotinylated Anti-Murine IL-22 Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mIL-22. Anti-Murine IL-22 specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot

To detect mIL-22 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIL-22 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect mIL-22 by sandwich ELISA (using $100 \, \mu$ l/well antibody solution) a concentration of $0.25 - 1.0 \, \mu$ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti-Murine IL-22 (61-022P) as a capture antibody, allows the detection of at least $0.2 - 0.4 \, n$ g/well of recombinant mIL-22.

Direct

To detect mIL-22 by direct ELISA (using 100 μ l/well antibody solution) a concentration of 0.25 – 1.0 μ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant mIL-22.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Biotinylated Anti-Murine IL-22 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Biotinylated Anti-Murine IL-22 Antibody - Protocols





Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Biotinylated Anti-Murine IL-22 Antibody - Images