

Biotinylated Anti-Rat IL-7 Antibody
Catalog # ABG10306**Specification**

Biotinylated Anti-Rat IL-7 Antibody - Product Information

Application	WB, E
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal

Biotinylated Anti-Rat IL-7 Antibody - Additional Information**Preparation**

Produced from sera of rabbits immunized with highly pure recombinant Rat IL-7. Anti-Rat IL-7 specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot

To detect Rat IL-7 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 mg/ml. When used in conjunction with compatible development reagents the detection limit for recombinant Rat IL-7 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Rat IL-7 by sandwich ELISA (using 100 ml/well) a concentration of 0.25 - 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems's Polyclonal Anti-Rat IL-7 () as a capture antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IL-7.

Direct

To detect Rat IL-7 by direct ELISA (using 100 ml/well) a concentration of approximately 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IL-7.

Storage

-20°C

Precautions

Biotinylated Anti-Rat IL-7 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Biotinylated Anti-Rat IL-7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Biotinylated Anti-Rat IL-7 Antibody - Images