

Anti-Human IL-8 Antibody
Catalog # ABG10310**Specification**

Anti-Human IL-8 Antibody - Product Information

Application	WB, E
Reactivity	Human
Host	Mouse
Clonality	Monoclonal

Anti-Human IL-8 Antibody - Additional Information**Preparation**

Produced in mice using highly pure (>98%) recombinant human IL-8 as the immunizing antigen. This IgG1 antibody was purified from cell culture by Protein G affinity chromatography.

WesternBlot

To detect hIL-8 by Western Blot analysis this antibody can be used at a concentration of 0.20-0.40 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hIL-8 is 2.0-4.0 ng/lane, under non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming 100µl/well), a concentration of 2.0-4.0 µg/ml of this antibody will detect at least 1000 pg/ml of recombinant human IL-8 when used with BioGems's biotinylated antigen affinity purified anti-human IL-8 (60-008BT) as the detection antibody at a concentration of approximately 0.5-1.0 µg/ml.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human IL-8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human IL-8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human IL-8 Antibody - Images