

Anti-Human IL-9 Antibody

Catalog # ABG10311

Specification

Anti-Human IL-9 Antibody - Product Information

Application WB, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human IL-9 Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure recombinant Human IL-9. Anti-Human IL-9 specific antibody was purified by affinity chromatography employing immobilized Human IL-9 matrix.

WesternBlot

To detect Human IL-9 by Western Blot analysis this antibody can be used at a concentration of $0.1 - 0.2\mu g/ml$. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human IL-9 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Human IL-9 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human IL-9 (60-009BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human IL-9.

Neutralization

To yield one-half maximal inhibition [ND_{>50}] of the biological activity of Human IL-9 (5.0 ng/ml), a concentration of 1.0-2.0 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human IL-9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human IL-9 Antibody - Protocols







Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human IL-9 Antibody - Images