

Anti-Rat IP-10 Antibody

Catalog # ABG10317

## Specification

# Anti-Rat IP-10 Antibody - Product Information

Application	WB, IHC, E
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal

# **Anti-Rat IP-10 Antibody - Additional Information**

#### Preparation

Produced from sera of rabbits immunized with highly pure recombinant Rat IP-10. Anti-Rat IP-10 specific antibody was purified by affinity chromatography employing an immobilized Rat IP-10 matrix.

### WesternBlot

To detect Rat IP-10 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2  $\mu$ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat IP-10 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

## Sandwich

To detect Rat IP-10 by sandwich ELISA (using 100 $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Rat IP-10 (62-062BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IP-10.

#### Immunohistochemistry

This antibody stained colchicine injected rat brain (including the caudate putamen) tissue. The primary antibody was incubated at 0.25 µg/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA<sup>™</sup>) reagent. Optimal concentrations and conditions may vary. <strong><em><span style="font-size: 10px;">Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm. </span></em>

#### Neutralization

To yield one-half maximal inhibition [<strong>ND</strong><span style="font-size: 16px;"><sub>50</sub></span>] of the biological activity of Rat IP-10 (100 ng/ml), a concentration of 5.0-10.0  $\mu$ g/ml of this antibody is required.

# **Formulation** A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

# Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage -20°C



Precautions

Anti-Rat IP-10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# **Anti-Rat IP-10 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Rat IP-10 Antibody - Images