10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Biotinylated Anti-Murine IP-10 Antibody
Catalog # ABG10319

Specification

Biotinylated Anti-Murine IP-10 Antibody - Product Information

Application	**WB, E**
Reactivity	**Mouse**
Host	**Rabbit**
Clonality	**Polyclonal**

Biotinylated Anti-Murine IP-10 Antibody - Additional Information

Preparation
Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mIP-10. Anti-Murine IP-10 specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot
To detect mIP-10 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIP-10 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich
To detect mIP-10 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.25 – 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti-Murine IP-10 (61-081P) as a capture antibody, allows the detection of at least 0.2 – 0.4 ng/well of recombinant mIP-10.

Direct
To detect mIP-10 by direct ELISA (using 100 µl/well antibody solution) a concentration of 0.25 – 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant mIP-10.

Formulation
A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution
Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage
-20°C

Precautions
Biotinylated Anti-Murine IP-10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Biotinylated Anti-Murine IP-10 Antibody - Protocols

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Biotinylated Anti-Murine IP-10 Antibody - Images