

Anti-Human Klotho Antibody

Catalog # ABG10332

Specification

Anti-Human Klotho Antibody - Product Information

Application WB, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human Klotho Antibody - Additional Information

Preparation

<span id="ctl00_SPWebPartManager1_g_2c64f2a8_025c_4a21_8444_332def32454d_ctl00_lblPrep
arationText">Produced from sera of rabbits pre-immunized with highly pure recombinant Human
Klotho. Anti-Human Klotho specific antibody was purified by affinity chromatography employing
immobilized Human Klotho matrix./span>

WesternBlot

To detect hKlotho by Western Blot analysis this antibody can be used at a concentration of 0.1 - $0.2~\mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hKlotho is 1.5 - 3.0~ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hKlotho by sandwich ELISA (using 100μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human Klotho () as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hKlotho./span>

Storage -20°C

Precautions

Anti-Human Klotho Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human Klotho Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation





Flow CytometyCell Culture

Anti-Human Klotho Antibody - Images