

## Anti-Rat MCP-1 Antibody

Catalog # ABG10353

# **Specification**

## **Anti-Rat MCP-1 Antibody - Product Information**

Application WB, IHC, E
Reactivity Rat
Host Rabbit
Clonality Polyclonal

## **Anti-Rat MCP-1 Antibody - Additional Information**

#### **Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant Rat MCP-1(MCAF). Anti-Rat MCP-1(MCAF) specific antibody was purified by affinity chromatography employing immobilized Rat MCP-1(MCAF) matrix.

#### WesternBlot

To detect Rat MCP-1(MCAF) by Western Blot analysis this antibody can be used at a concentration of 0.1 -  $0.2~\mu g/ml$ . Used in conjunction with compatible secondary reagents the detection limit for recombinant Rat MCP-1(MCAF) is 1.5-3.0~ng/lane, under either reducing or non-reducing conditions.

#### Sandwich

To detect Rat MCP-1(MCAF) by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Rat MCP-1(MCAF) (62-065BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat MCP-1(MCAF).

#### **Immunohistochemistry**

This antibody stained colchicine injected rat brain (including the caudate putamen) tissue. The primary antibody was incubated at 1.0  $\mu$ g/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA $^{\text{TM}}$ ) reagent. Optimal concentrations and conditions may vary. <strong><span style="font-size: 10px;"><em>Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm. </em></strong>

#### **Neutralization**

To yield one-half maximal inhibition [<strong>ND</strong><span style="font-size: 16px;"><sub>50</sub></span>] of the biological activity of Rat MCP-1(MCAF)(100 ng/ml), a concentration of 4.0-6.0 µg/ml of this antibody is required.

#### **Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

#### Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

# Storage -20°C



# **Precautions**

Anti-Rat MCP-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# **Anti-Rat MCP-1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

**Anti-Rat MCP-1 Antibody - Images**