

# **Biotinylated Anti-Human MCP-1 Antibody**

Catalog # ABG10354

# **Specification**

# Biotinylated Anti-Human MCP-1 Antibody - Product Information

Application WB, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

## Biotinylated Anti-Human MCP-1 Antibody - Additional Information

#### **Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMCP-1/MCAF. Anti-Human MCP-1/MCAF specific antibody was purified by affinity chromatography and then biotinylated.

#### WesternBlot

To detect hMCP-1/MCAF by Western Blot analysis this antibody can be used at a concentration of 0.1 -  $0.2~\mu g/ml$ . Used in conjunction with compatible secondary reagents the detection limit for recombinant hMCP-1/MCAF is 1.5 - 3.0~n g/lane, under either reducing or non-reducing conditions.

#### Sandwich

To detect hMCP-1/MCAF by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.25 – 1.0  $\mu$ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti-Human MCP-1/MCAF (60-212P) as a capture antibody, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hMCP-1/MCAF.

## **Direct**

To detect hMCP-1/MCAF by direct ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.25 – 1.0  $\mu$ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hMCP-1/MCAF.

#### **Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

#### Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

#### **Storage**

-20°C

## **Precautions**

Biotinylated Anti-Human MCP-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



# **Biotinylated Anti-Human MCP-1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**Biotinylated Anti-Human MCP-1 Antibody - Images**