

# **Biotinylated Anti-Human MCP-3 Antibody**

Catalog # ABG10365

## Specification

## **Biotinylated Anti-Human MCP-3 Antibody - Product Information**

Application Reactivity Host Clonality WB, E Human Goat Polyclonal

## **Biotinylated Anti-Human MCP-3 Antibody - Additional Information**

### Preparation

Produced from sera of Goats pre-immunized with highly pure (>98%) recombinant hMCP-3. Anti-Human MCP-3 specific antibody was purified by affinity chromatography and then biotinylated.

#### WesternBlot

To detect by Western Blot analysis this antibody can be used at a concentration of  $0.1 - 0.2 \mu g/ml$ . Used in conjunction with compatible secondary reagents the detection limit for recombinant hMCP-3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

### Sandwich

To detect hMCP-3 by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.25 – 1.0  $\mu$ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti-Human MCP-3 (60-214GP) as a capture antibody, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hMCP-3.

#### Direct

To detect hMCP-3 by direct ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.25 – 1.0  $\mu$ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hMCP-3.

## Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

#### Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage -20°C

#### Precautions

Biotinylated Anti-Human MCP-3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



## **Biotinylated Anti-Human MCP-3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

**Biotinylated Anti-Human MCP-3 Antibody - Images**