

Anti-Human MCP-4 Antibody
Catalog # ABG10373**Specification**

Anti-Human MCP-4 Antibody - Product Information

Application	WB, E
Reactivity	Human
Host	Mouse
Clonality	Monoclonal

Anti-Human MCP-4 Antibody - Additional Information**Preparation**

Produced in BALB/c mice using highly pure (>98%) recombinant human MCP-4 as the immunizing antigen. This IgG1_K antibody was purified from cell culture by Protein A affinity chromatography.

WesternBlot

To detect hMCP-4 by Western Blot analysis this antibody can be used at a concentration of 0.20-0.40 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMCP-4 is 2.0-4.0 ng/lane, under non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming 100µl/well), a concentration of 2.0-4.0 µg/ml of this antibody will detect at least 100 pg/ml of recombinant human MCP-4 when used with BioGems's biotinylated antigen affinity purified anti-human MCP-4 (60-215BT) as the detection antibody at a concentration of approximately 0.5-1.0 µg/ml.

Neutralization

To yield one-half maximal inhibition [ND₅₀] of the biological activity of Human MCP-4 (100 ng/ml), a concentration of 0.8-1.0 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human MCP-4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human MCP-4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human MCP-4 Antibody - Images