

Anti-Human MEC Antibody

Catalog # ABG10385

Specification

Anti-Human MEC Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human MEC Antibody - Additional Information

Preparation

Produced from sera of rabbits immunized with highly pure recombinant Human MEC. Anti-Human MEC specific antibody was purified by affinity chromatography employing an immobilized Human MEC matrix.

WesternBlot

To detect hMEC by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 $\mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMEC is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hMEC by sandwich ELISA (using 100μ I/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human MEC (60-218BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMEC.

Immunohistochemistry

This antibody stained CACO-2 cells. The primary antibody was incubated at 3.0 μ g/ml overnight at 4°C followed by a fluorescent labeled secondary antibody. Optimal concentrations and conditions may vary. Information courtesy of the Cell Profiling group, SciLifeLab Stockholm.

sciLifeLab Stockholm.<a href="mailto:sciLifeLab Stockholm.<a href="mailto:sciLifeLab Stockholm.<a href="mailto:sciLifeLab Stockholm.<a href="mailto:sciLifeLab Stockholm.<a href="mailto:sciLifeLab Stockholm.<a href="mailto:sciLifeLab Stockholm.<a href="mailto:sciL

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human MEC Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



Anti-Human MEC Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human MEC Antibody - Images