

Anti-Human MIP-1 β Antibody
Catalog # ABG10405**Specification**

Anti-Human MIP-1 β Antibody - Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Anti-Human MIP-1 β Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure recombinant Human MIP-1 β . Anti-Human MIP-1 β specific antibody was purified by affinity chromatography employing immobilized Human MIP-1 β matrix.

WesternBlot

To detect Human MIP-1 β by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human MIP-1 β is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Human MIP-1 β by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human MIP-1 β (60-225BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human MIP-1 β .

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human pancreas adenocarcinoma. The recommended concentration is 1.0 mg/ml with 2 hour incubation at room temperature. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

Neutralization

To yield one-half maximal inhibition [**ND**">₅₀] of the biological activity of Human MIP-1 β (50.00 ng/ml), a concentration of 1.75-3.00 μ g/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human MIP-1 β Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human MIP-1 β Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human MIP-1 β Antibody - Images