

Anti-Human MIP-3α Antibody

Catalog # ABG10415

Specification

Anti-Human MIP-3α Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human MIP-3α Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure recombinant Human MIP-3 α . Anti-Human MIP-3 α specific antibody was purified by affinity chromatography employing immobilized Human MIP-3 α matrix.

WesternBlot

To detect HumanMIP-3 α by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human MIP-3 α is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Human MIP-3 α by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human MIP-3 α (60-227BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human MIP-3 α .

Immunohistochemistry

This antibody stained formalin-fixed paraffin-embedded sections of human colon/rectum adenocarcinoma tissue. The recommended concentration is $0.5\mu g/ml$ with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.


Storage

-20°C

Precautions

Anti-Human MIP- 3α Antibody is for research use only and not for use in diagnostic or therapeutic procedures.







Anti-Human MIP-3α Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human MIP-3α Antibody - Images