

**Anti-Human NAP-2 Antibody**  
**Catalog # ABG10431****Specification**

---

**Anti-Human NAP-2 Antibody - Product Information**

Application	<b>WB, E</b>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>

**Anti-Human NAP-2 Antibody - Additional Information****Preparation**

Produced in BALB/c mice using highly pure (>98%) recombinant human NAP-2 as the immunizing antigen. This IgG1<sub>K</sub> antibody was purified from cell culture by ammonium sulfate co-precipitation.

**WesternBlot**

To detect hNAP-2 by Western Blot analysis this antibody can be used at a concentration of 0.5-1.0 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNAP-2 is 0.5-1.0 ng/lane, under reducing or non-reducing conditions.

**Sandwich**

In a sandwich ELISA (assuming 100µl/well), a concentration of 2.0-4.0 µg/ml of this antibody will detect at least 500 pg/ml of recombinant human NAP-2 when used with BioGems's biotinylated antigen affinity purified anti-human NAP-2 (60-242BT) as the detection antibody at a concentration of approximately 0.5-1.0 µg/ml.

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS.

**Reconstitution**

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Human NAP-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-Human NAP-2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-Human NAP-2 Antibody - Images**