

Anti-Human NP-1 Antibody

Catalog # ABG10440

Specification

Anti-Human NP-1 Antibody - Product Information

Application Reactivity Host Clonality WB, E Human Goat Polyclonal

Anti-Human NP-1 Antibody - Additional Information

Preparation

Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hNP-1. Anti-Human NP-1 specific antibody was purified by affinity chromatography employing immobilized hNP-1 matrix.

WesternBlot

To detect hNP-1 by Western Blot analysis this antibody can be used at a concentration of 0.1- 0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNP-1 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hNP-1 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human NP-1 (60-251BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hNP-1.

Neutralization

To yield one-half maximal inhibition [ND₅₀] of the biological activity of Human NP-1 (100.0 ng/ml), a concentration of 0.40-0.60 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage -20°C

Precautions

Anti-Human NP-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human NP-1 Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Human NP-1 Antibody - Images