

# **Anti-Human Resistin Antibody**

Catalog # ABG10484

# **Specification**

# **Anti-Human Resistin Antibody - Product Information**

Application WB, IHC, E
Reactivity Human
Host Mouse
Clonality Monoclonal

# **Anti-Human Resistin Antibody - Additional Information**

#### **Preparation**

Produced in BALB/c mice using highly pure (>98%) recombinant human Resistin as the immunizing antigen. This  $IgG < sub > 2b\kappa < / sub >$  antibody was purified from cell culture by Protein A affinity chromatography.

#### WesternBlot

To detect hResistin by Western Blot analysis this antibody can be used at a concentration of 0.20- $0.40 \mu g/ml$ . Used in conjunction with compatible secondary reagents the detection limit for recombinant hResistin is 0.20- $0.40 \mu g/lane$ , under non-reducing conditions.

### Sandwich

In a sandwich ELISA (assuming  $100\mu$ I/well), a concentration of 4.0- $8.0~\mu$ g/ml of this antibody will detect at least 200~pg/ml of recombinant human Resistin when used with BioGems' biotinylated antigen affinity purified anti-human Resistin (60-289BT) as the detection antibody at a concentration of at least 0.5- $1.0~\mu$ g/ml.

## **Immunohistochemistry**

This antibody stained formalin-fixed, paraffin-embedded sections of human normal placenta. The recommended concentration is  $1.0~\mu g/ml$  with a two hour incubation at room temperature. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

#### **Formulation**

A sterile filtered antibody solution was lyophilized from PBS.

# Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

# **Storage**

-20°C

### **Precautions**

Anti-Human Resistin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



# **Anti-Human Resistin Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**Anti-Human Resistin Antibody - Images**