

Anti-Murine SDF-1α Antibody

Catalog # ABG10501

Specification

Anti-Murine SDF-1α Antibody - Product Information

Application	WB, E
Reactivity	Mouse
Host	Goat
Clonality	Polyclonal

Anti-Murine SDF-1a Antibody - Additional Information

Preparation

Produced from sera of goats pre-immunized with highly pure recombinant Murine SDF-1 α . Anti-Murine SDF-1 α specific antibody was purified by affinity chromatography employing immobilized Murine SDF-1 α matrix.

WesternBlot

To detect Murine SDF-1 α by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Murine SDF-1 α is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Murine SDF-1 α by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Murine SDF-1 α (61-120BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Murine SDF-1 α .

Neutralization

To yield one-half maximal inhibition [ND₅₀] of the biological activity of Murine SDF-1 α (100 ng/ml), a concentration of 2.5 - 3.5 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage -20°C

Precautions

Anti-Murine SDF-1 α Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Murine SDF-1a Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Murine SDF-1a Antibody - Images