

Anti-Human sFas Receptor Antibody
Catalog # ABG10510**Specification**

Anti-Human sFas Receptor Antibody - Product Information

Application	WB, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Anti-Human sFas Receptor Antibody - Additional Information**Preparation**

Produced from sera of rabbits immunized with highly pure recombinant Human sFas Receptor. Anti-Human sFas Receptor specific antibody was purified by affinity chromatography employing an immobilized Human sFas Receptor matrix.

WesternBlot

To detect Human sFas Receptor by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. When used in conjunction with compatible secondary reagents the detection limit for recombinant Human sFas Receptor is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Human sFas Receptor by sandwich ELISA (using 100µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human sFas Receptor (60-130BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Human sFas Receptor.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human sFas Receptor Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human sFas Receptor Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human sFas Receptor Antibody - Images