

**Anti-Murine TNF- $\alpha$  Antibody**  
**Catalog # ABG10547****Specification****Anti-Murine TNF- $\alpha$  Antibody - Product Information**

Application	WB, IHC, E
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal

**Anti-Murine TNF- $\alpha$  Antibody - Additional Information****Preparation**

Produced from sera of rabbits pre-immunized with highly pure recombinant Murine TNF- $\alpha$ . Anti-Murine TNF- $\alpha$  specific antibody was purified by affinity chromatography employing immobilized Murine TNF- $\alpha$  matrix.

**WesternBlot**

To detect Murine TNF- $\alpha$  by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2  $\mu$ g/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Murine TNF- $\alpha$  is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

**Sandwich**

To detect Murine TNF- $\alpha$  by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Murine TNF- $\alpha$  (61-124BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Murine TNF- $\alpha$ .

**Immunohistochemistry**

This antibody stained colchicine injected mouse brain (hippocampus CA1 region) tissue. The primary antibody was incubated at 1.0  $\mu$ g/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA™) reagent. Optimal concentrations and conditions may vary.   
Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm.

**Neutralization**

To yield one-half maximal inhibition [ $\text{ND}_{50}$ ] of the biological activity of Murine TNF- $\alpha$  (0.5 ng/ml), a concentration of 0.04 - 0.05  $\mu$ g/ml of this antibody is required.

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Reconstitution**

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Murine TNF- $\alpha$  Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-Murine TNF- $\alpha$  Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-Murine TNF- $\alpha$  Antibody - Images**