10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Human TWEAK Antibody
Catalog # ABG10569

Specification

Anti-Human TWEAK Antibody - Product Information

Application	**WB, E**
Reactivity	**Human**
Host	**Goat**
Clonality	**Polyclonal**

Anti-Human TWEAK Antibody - Additional Information

Preparation
Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hTWEAK. Anti-Human TWEAK specific antibody was purified by affinity chromatography employing immobilized hTWEAK matrix.

WesternBlot
To detect hTWEAK by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hTWEAK is 1.5-3.0 ng/lane, under either reducing or or non-reducing conditions.

Sandwich
To detect hTWEAK by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human TWEAK (60-335BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hTWEAK.

Neutralization
To yield one-half maximal inhibition [ND50] of the biological activity of hTWEAK (200 ng/ml), a concentration of 0.1 - 0.2 µg/ml of this antibody is required.

Formulation
A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution
Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage
-20°C

Precautions
Anti-Human TWEAK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human TWEAK Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human TWEAK Antibody - Images