10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-SMAD1/SMAD5 Picoband Antibody

Catalog # ABO10105

Specification

Anti-SMAD1/SMAD5 Picoband Antibody - Product Information

Application	**WB**
Primary Accession	Q15797
Host	**Rabbit**
Reactivity	**Human, Mouse, Rat**
Clonality	**Polyclonal**
Format	**Lyophilized**

Description
Rabbit IgG polyclonal antibody for SMAD1/SMAD5 detection. Tested with WB in Human;Mouse;Rat.

Reconstitution
Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-SMAD1/SMAD5 Picoband Antibody - Additional Information

Gene ID 4086

Other Names
Mothers against decapentaplegic homolog 1, MAD homolog 1, Mothers against DPP homolog 1, JV4-1, Mad-related protein 1, SMAD family member 1, SMAD 1, Smad1, hSMAD1, Transforming growth factor-beta-signaling protein 1, BSP-1, SMAD1, BSP1, MADH1, MADR1

Application Details
Western blot, 0.1-0.5 µg/ml

Subcellular Localization
Cytoplasm.

Tissue Specificity
Ubiquitous. Highest expression seen in the heart and skeletal muscle.

Contents
Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen
A synthetic peptide corresponding to a sequence of human SMAD1/SMAD5 (KRLLGWKQGDEEEKWAEKAVDALVKKLKKKKGAMEELEK).

Cross Reactivity
No cross reactivity with other proteins.

Storage	**At -20˚C; for one year. After r˚Constitution, at 4˚C; for one month. It˚Can also be aliquotted and stored frozen at -20˚C; for a**

longer time. Avoid repeated freezing and thawing.

Anti-SMAD1/SMAD5 Picoband Antibody - Protein Information

Name SMAD1

Synonyms BSP1, MADH1, MADR1

Function
Transcriptional modulator that plays a role in various cellular processes, including embryonic development, cell differentiation, and tissue homeostasis (PubMed:9335504). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRIs) and associates with SMAD4 to form an heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed:33667543). In turn, the hetero-trimeric complex recognizes cis-regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed:33667543). SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1. Positively regulates BMP4-induced expression of odontogenic development regulator MSX1 following IPO7-mediated nuclear import (By similarity).

Cellular Location
Cytoplasm. Nucleus Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4 (PubMed:15647271). Co-localizes with LEMD3 at the nucleus inner membrane (PubMed:15647271). Exported from the nucleus to the cytoplasm when dephosphorylated (By similarity) {ECO:0000250|UniProtKB:P70340, ECO:0000269|PubMed:15647271}

Tissue Location
Ubiquitous. Highest expression seen in the heart and skeletal muscle

Anti-SMAD1/SMAD5 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-SMAD1/SMAD5 Picoband Antibody - Images