10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-DCK Picoband Antibody
Catalog # ABO10203

Specification

Anti-DCK Picoband Antibody - Product Information

Application	**WB, E**
Primary Accession	A01655-1
Host	**Rabbit**
Reactivity	**Human, Mouse, Rat**
Clonality	**Polyclonal**
Format	**Lyophilized**

Description
Rabbit IgG polyclonal antibody for DCK detection. Tested with WB, Direct ELISA in Human;Mouse;Rat.

Reconstitution
Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-DCK Picoband Antibody - Additional Information

Application Details
Western blot, 0.1-0.5 µg/ml
 Direct ELISA, 0.1-0.5 µg/ml

Subcellular Localization
Nucleus.

Contents
Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen
E. coli-derived human DCK recombinant protein (Position: E17-L260).

Cross Reactivity
No cross reactivity with other proteins.

Storage	**At -20˚C; for one year. After r˚Constitution, at 4˚C; for one month. It˚Can also be aliquotted and stored frozen at -20˚C; for a longer time. Avoid repeated freezing and thawing.**

Anti-DCK Picoband Antibody - Protein Information

Anti-DCK Picoband Antibody - Protocols

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-DCK Picoband Antibody - Images

Anti-DCK Picoband Antibody - Background

Deoxycytidine kinase (dCK) is an enzyme which is encoded by the DCK gene in humans. Deoxycytidine kinase (DCK) is required for the phosphorylation of several deoxyribonucleosides and their nucleoside analogs. Deficiency of DCK is associated with resistance to antiviral and anticancer chemotherapeutic agents. Conversely, increased deoxycytidine kinase activity is associated with increased activation of these compounds to cytotoxic nucleoside triphosphate derivatives. DCK is clinically important because of its relationship to drug resistance and sensitivity.