

**Anti-NMDAR1 Antibody**  
**Catalog # ABO10600****Specification**

---

**Anti-NMDAR1 Antibody - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">Q05586</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Glutamate receptor ionotropic, NMDA 1 (GRIN1) detection. Tested with WB, IHC-P in Human; Mouse; Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-NMDAR1 Antibody - Additional Information**

**Gene ID** 2902

**Other Names**

Glutamate receptor ionotropic, NMDA 1, GluN1, Glutamate [NMDA] receptor subunit zeta-1, N-methyl-D-aspartate receptor subunit NR1, NMD-R1, GRIN1, NMDAR1

**Calculated MW**

105373 MW KDa

**Application Details**

Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat  
Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse

**Subcellular Localization**

Cell membrane ; Multi-pass membrane protein . Cell junction, synapse, postsynaptic cell membrane . Cell junction, synapse, postsynaptic cell membrane, postsynaptic density . Enriched in postsynaptic plasma membrane and postsynaptic densities. .

**Protein Name**

Glutamate receptor ionotropic, NMDA 1

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human NMDAR1(36-53aa RKHEQMFREAVNQANKRH), identical to the related rat and mouse sequences.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Belongs to the glutamate-gated ion channel (TC 1.A.10.1) family. NR1/GRIN1 subfamily.

**Anti-NMDAR1 Antibody - Protein Information**

**Name** GRIN1

**Synonyms** NMDAR1

**Function**

Component of NMDA receptor complexes that function as heterotetrameric, ligand-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium. Channel activation requires binding of the neurotransmitter glutamate to the epsilon subunit, glycine binding to the zeta subunit, plus membrane depolarization to eliminate channel inhibition by Mg(2+) (PubMed: [7685113](http://www.uniprot.org/citations/7685113), PubMed: [28126851](http://www.uniprot.org/citations/28126851), PubMed: [26919761](http://www.uniprot.org/citations/26919761), PubMed: [26875626](http://www.uniprot.org/citations/26875626), PubMed: [28105280](http://www.uniprot.org/citations/28105280)). Sensitivity to glutamate and channel kinetics depend on the subunit composition (PubMed: [26919761](http://www.uniprot.org/citations/26919761)).


**Cellular Location**


Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane. Postsynaptic density. Note=Enriched in postsynaptic plasma membrane and postsynaptic densities.

**Anti-NMDAR1 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-NMDAR1 Antibody - Images**



Anti-NMDAR1 antibody, ABO10600, Western blotting  
Lane 1: Rat Brain Tissue Lysate  
Lane 2: Rat Brain Tissue Lysate  
Lane 3: Rat Liver Tissue Lysate  
Lane 4: Rat Heart Tissue Lysate  
Lane 5: MM453 Cell Lysate  
Lane 6: MM231 Cell Lysate  
Lane 7: HELA Cell Lysate



Anti-NMDAR1 antibody, ABO10600, IHC(P)  
IHC(P): Rat Brain Tissue

### **Anti-NMDAR1 Antibody - Background**

Glutamate [NMDA] receptor subunit zeta-1 is a protein that in humans is encoded by the GRIN1 gene. The protein encoded by this gene is a critical subunit of N-methyl-D-aspartate receptors, members of the glutamate receptor channel superfamily which are heteromeric protein complexes with multiple subunits arranged to form a ligand-gated ion channel. These subunits play a key role in the plasticity of synapses, which is believed to underlie memory and learning. Cell-specific factors are thought to control expression of different isoforms, possibly contributing to the functional diversity of the subunits. Alternatively spliced transcript variants have been described.