

Anti-FGF19 Picoband Antibody
Catalog # ABO11680**Specification**

Anti-FGF19 Picoband Antibody - Product Information

Application	WB
Primary Accession	Q8VI81
Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Fibroblast growth factor 15(FGF19) detection. Tested with WB in Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-FGF19 Picoband Antibody - Additional Information**Calculated MW**

25207 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Rat

Protein Name

Fibroblast growth factor 15

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived rat FGF19 recombinant protein (Position: R26-K218). Rat FGF19 shares 53.2% and 96.4% amino acid (aa) sequence identity with human and mouse FGF19, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.


Anti-FGF19 Picoband Antibody - Protein Information

Anti-FGF19 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-FGF19 Picoband Antibody - Images



Western blot analysis of FGF19 expression in rat pancreas extract (lane 1) and NRK whole cell lysates (lane 2). FGF19 at 24KD was detected using rabbit anti- FGF19 Antigen Affinity purified polyclonal antibody (Catalog # ABO11680) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .

Anti-FGF19 Picoband Antibody - Background

FGF19, Fibroblast growth factor 19, is a protein that in humans is encoded by the FGF19 gene. The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. The FGF19 gene is mapped to 11q13.3. The deduced 216-amino acid FGF19 protein contains a signal sequence and 2 cysteine residues that are conserved in the FGF family. Expression of this gene was detected only in fetal but not adult brain tissue. Synergistic interaction of the chick homolog and Wnt-8c has been shown to be required for initiation of inner ear development. FGF19 stimulates hepatic protein and glycogen synthesis but does not induce lipogenesis. The effects of FGF19 are independent of the activity of either insulin or the protein kinase Akt and, instead, are mediated through a mitogen-activated protein kinase signaling pathway that activates components of the protein translation machinery and stimulates glycogen synthase activity. The orthologous protein in mouse is FGF15, which shares about 50% amino acid identity and has similar functions. Together they are often referred to as FGF15/19.