

Anti-PINX1 Picoband Antibody
Catalog # ABO12186**Specification**

Anti-PINX1 Picoband Antibody - Product Information

Application	WB
Primary Accession	Q96BK5
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for PIN2/TERF1-interacting telomerase inhibitor 1(PINX1) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-PINX1 Picoband Antibody - Additional Information

Gene ID 54984

Other Names

PIN2/TERF1-interacting telomerase inhibitor 1, Liver-related putative tumor suppressor, Pin2-interacting protein X1, Protein 67-11-3, TRF1-interacting protein 1, PINX1, LPTL, LPTS

Calculated MW

37035 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Nucleus. Nucleus, nucleolus. Chromosome, telomere. Chromosome, centromere, kinetochore. Localizes in nucleoli, at telomere speckles and to the outer plate of kinetochores. Localization to the kinetochore is mediated by its central region and depends on NDC80 and CENPE.

Tissue Specificity

Ubiquitous; expressed at low levels. Not detectable in a number of hepatocarcinoma cell lines.

Protein Name

PIN2/TERF1-interacting telomerase inhibitor 1

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human PINX1 (54-83aa DHIKVQVKNNHLGLGATINNEDNWIAHQDD), different from the related mouse and rat sequences by

three amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the PINX1 family.

Anti-PINX1 Picoband Antibody - Protein Information

Name PINX1

Synonyms LPTL, LPTS

Function

Microtubule-binding protein essential for faithful chromosome segregation. Mediates TRF1 and TERT accumulation in nucleolus and enhances TRF1 binding to telomeres. Inhibits telomerase activity. May inhibit cell proliferation and act as tumor suppressor.

Cellular Location

Nucleus. Nucleus, nucleolus. Chromosome, telomere. Chromosome, centromere, kinetochore
Note=Localizes in nucleoli, at telomere speckles and to the outer plate of kinetochores.
Localization to the kinetochore is mediated by its central region and depends on NDC80 and CENPE


Tissue Location


Ubiquitous; expressed at low levels. Not detectable in a number of hepatocarcinoma cell lines

Anti-PINX1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PINX1 Picoband Antibody - Images



Anti- PINX1 Picoband antibody, ABO12186, Western blotting All lanes: Anti PINX1 (ABO12186) at 0.5ug/ml
Lane 1: Human Placenta Tissue Lysate at 50ug
Lane 2: HELA Whole Cell Lysate at 40ug
Lane 3: HUT Whole Cell Lysate at 40ug
Lane 4: JURKAT Whole Cell Lysate at 40ug
Predicted bind size: 37KD
Observed bind size: 37KD

Anti-PINX1 Picoband Antibody - Background

PINX1, also known as PIN2 interacting protein 1, is a telomerase inhibitor and a possible tumor suppressor. It is mapped to 8p23. Over-expression of PINX1 results in decreased telomerase activity, telomere shortening, and induction of crisis. Reduction of PINX1 leads to an increase in telomerase activity and elongation of telomeres. PINX1 differs from other proteins that regulate telomere length in that it acts on telomerase while other proteins adjust telomere length without affecting telomerase activity. The PINX1 budding yeast orthologue Gnop1 inhibits telomerase by isolating the uncomplexed TERT protein so that it cannot associate with the telomerase template RNA, which prevents telomerase from being assembled. However, in humans, PINX1 impedes already assembled telomerase.