

Anti-Fascin Picoband Antibody
Catalog # ABO12278**Specification**

Anti-Fascin Picoband Antibody - Product Information

Application	WB
Primary Accession	Q16658
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Fascin(FSCN1) detection. Tested with WB in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Fascin Picoband Antibody - Additional Information

Gene ID 6624

Other Names

Fascin, 55 kDa actin-bundling protein, Singed-like protein, p55, FSCN1, FAN1, HSN, SNL

Calculated MW

54530 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Rat

Subcellular Localization

Cytoplasm, cytoskeleton. Cell projection, filopodium. Cell projection, invadopodium. Cytoplasm, cytosol. In glioma cells, partially colocalizes with F-actin stress fibers in the cytosol.

Tissue Specificity

Ubiquitous.

Protein Name

Fascin

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human Fascin (42-73aa KKQIWTLEQPPDEAGSAAVCLRSHLGRYLAAD), identical to the related mouse sequence, and different from the related rat sequence by one amino acid.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the fascin family.

Anti-Fascin Picoband Antibody - Protein Information

Name FSCN1

Synonyms FAN1, HSN, SNL

Function

Actin-binding protein that contains 2 major actin binding sites (PubMed: [21685497](http://www.uniprot.org/citations/21685497), PubMed: [23184945](http://www.uniprot.org/citations/23184945)). Organizes filamentous actin into parallel bundles (PubMed: [20393565](http://www.uniprot.org/citations/20393565), PubMed: [21685497](http://www.uniprot.org/citations/21685497), PubMed: [23184945](http://www.uniprot.org/citations/23184945)). Plays a role in the organization of actin filament bundles and the formation of microspikes, membrane ruffles, and stress fibers (PubMed: [22155786](http://www.uniprot.org/citations/22155786)). Important for the formation of a diverse set of cell protrusions, such as filopodia, and for cell motility and migration (PubMed: [20393565](http://www.uniprot.org/citations/20393565), PubMed: [21685497](http://www.uniprot.org/citations/21685497), PubMed: [23184945](http://www.uniprot.org/citations/23184945)). Mediates reorganization of the actin cytoskeleton and axon growth cone collapse in response to NGF (PubMed: [22155786](http://www.uniprot.org/citations/22155786)).

Cellular Location

Cytoplasm, cytosol. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, stress fiber. Cell projection, filopodium. Cell projection, invadopodium. Cell projection, microvillus. Cell junction. Note=Colocalized with RUFY3 and F-actin at filopodia of the axonal growth cone. Colocalized with DBN1 and F-actin at the transitional domain of the axonal growth cone (By similarity). {ECO:0000250|UniProtKB:Q61553, ECO:0000269|PubMed:21706053}

Tissue Location

Ubiquitous.


Anti-Fascin Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Fascin Picoband Antibody - Images



Anti- Fascin Picoband antibody, ABO12278, Western blotting All lanes: Anti Fascin (ABO12278) at 0.5ug/ml
Lane 1: Rat Brain Tissue Lysate at 50ug
Lane 2: Rat Lung Tissue Lysate at 50ug
Lane 3: HELA Whole Cell Lysate at 40ug
Lane 4: SKOV Whole Cell Lysate at 40ug
Predicted bind size: 55KD
Observed bind size: 55KD

Anti-Fascin Picoband Antibody - Background

Fascin is an actin cross-linking protein. The Fascin gene contains 5 exons and spans 7 kb. It is a 54-58 kilodalton monomeric actin filament bundling protein originally isolated from sea urchin egg but also found in *Drosophila* and vertebrates, including humans. Fascin (from the Latin for bundle) is spaced at 11 nanometre intervals along the filament. The bundles in cross section are seen to be hexagonally packed, and the longitudinal spacing is compatible with a model where fascin cross-links at alternating 4 and 5 actins. It is calcium insensitive and monomeric. Fascin binds beta-catenin, and colocalizes with it at the leading edges and borders of epithelial and endothelial cells. The role of Fascin in regulating cytoskeletal structures for the maintenance of cell adhesion, coordinating motility and invasion through interactions with signalling pathways is an active area of research especially from the cancer biology perspective. Abnormal fascin expression or function has been implicated in breast cancer, colon cancer, esophageal squamous cell carcinoma, gallbladder cancer and prostate cancer.