

Anti-S100 Alpha 6 Picoband Antibody
Catalog # ABO12362**Specification**

Anti-S100 Alpha 6 Picoband Antibody - Product Information

Application	WB, IHC
Primary Accession	P06703
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Protein S100-A6(S100A6) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-S100 Alpha 6 Picoband Antibody - Additional Information

Gene ID 6277

Other Names

Protein S100-A6, Calcyclin, Growth factor-inducible protein 2A9, MLN 4, Prolactin receptor-associated protein, PRA, S100 calcium-binding protein A6, S100A6, CACY

Calculated MW

10180 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat

Western blot, 0.1-0.5 µg/ml, Human, Rat

Subcellular Localization

Nucleus envelope. Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side.

Protein Name

Protein S100-A6

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived human S100 alpha 6 recombinant protein (Position: M1-G90). Human S100 alpha 6 shares 96.6% and 95.5% amino acid (aa) sequence identity with mouse and rat S100 alpha 6, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-S100 Alpha 6 Picoband Antibody - Protein Information

Name S100A6

Synonyms CACY

Function

May function as calcium sensor and modulator, contributing to cellular calcium signaling. May function by interacting with other proteins, such as TPR-containing proteins, and indirectly play a role in many physiological processes such as the reorganization of the actin cytoskeleton and in cell motility. Binds 2 calcium ions. Calcium binding is cooperative.


Cellular Location

Nucleus envelope. Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side

Anti-S100 Alpha 6 Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-S100 Alpha 6 Picoband Antibody - Images




Anti- S100 alpha 6 Picoband antibody, ABO12362, Western blotting All lanes: Anti S100 alpha 6 (ABO12362) at 0.5ug/ml Lane 1: Rat Liver Tissue Lysate at 50ug Lane 2: A431 Whole Cell Lysate at 40ug Predicted bind size: 10KD Observed bind size: 10KD




Anti- S100 alpha 6 Picoband antibody, ABO12362, IHC(P) IHC(P): Mouse Testis Tissue



Anti- S100 alpha 6 Picoband antibody, ABO12362, IHC(P) IHC(P): Rat Lung Tissue



Anti- S100 alpha 6 Picoband antibody, ABO12362, IHC(P)IHC(P): Human Mammary Cancer Tissue



Anti- S100 alpha 6 Picoband antibody, ABO12362, IHC(P)IHC(P): Human Placenta Tissue

Anti-S100 Alpha 6 Picoband Antibody - Background

S100 calcium-binding protein A6 (S100A6) is a protein that in humans is encoded by the S100A6 gene. The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. And S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in stimulation of Ca^{2+} -dependent insulin release, stimulation of prolactin secretion, and exocytosis. Chromosomal rearrangements and altered expression of this gene have been implicated in melanoma.