

Bid Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10141**Specification**

Bid Antibody - Product Information

Application	WB, IHC, IP
Primary Accession	P55957
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21995

Bid Antibody - Additional Information**Gene ID 637**

Application & Usage	Western blot analysis (0.5-4 µg/ml), immunoprecipitation (5-10 µg/ml), and Immunohistochemistry (20-40 µg/ml). However, the optimal conditions should be determined individually. The antibody detects 22 kDa human Bid
---------------------	---

Other Names

FP497 , MGC42355, MGC15319

Target/Specificity

Bid

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) affinity purified rabbit anti-Bid polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

Bid Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Bid Antibody - Protein Information**Name** BID**Function**

Induces caspases and apoptosis (PubMed:14583606). Counters the protective effect of BCL2 (By similarity).

Cellular Location

Cytoplasm. Mitochondrion membrane. Mitochondrion outer membrane. Note=When uncleaved, it is predominantly cytoplasmic. [BH3-interacting domain death agonist p13]: Mitochondrion membrane {ECO:0000250|UniProtKB:P70444}. Note=Associated with the mitochondrial membrane. {ECO:0000250|UniProtKB:P70444} [Isoform 3]: Cytoplasm

Tissue Location

[Isoform 2]: Expressed in spleen, pancreas and placenta (at protein level). [Isoform 4]: Expressed in lung and pancreas (at protein level).

Bid Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bid Antibody - Images**Bid Antibody - Background**

Bid, a BH3 domain-containing proapoptotic Bcl-2 family member, is localized in the cytosolic fraction of cells as an inactive precursor. Its active form is generated upon proteolytic cleavage by caspase-8 in the Fas signaling pathway. Cleaved Bid translocates to mitochondria and releases its potent proapoptotic activity, which in turn induces cytochrome c release and mitochondrial damage. The cytochrome c releasing activity of Bid was antagonized by Bcl-2. Mutation in the SH3 domain can diminish the cytochrome c releasing activity. In the animal model studies, Bid-deficient mice are found resistant to the lethal effects of death factor signals relayed through Fas.