

FIAF Antibody

Rabbit Polyclonal Antibody Catalog # ABV10360

Specification

FIAF Antibody - Product Information

WB Application **Primary Accession** 09Z1P8 BAC25242 Other Accession Mouse, Rat Reactivity Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 45538

FIAF Antibody - Additional Information

Gene ID 57875

Application & Usage Western blotting (0.5-4 μg/ml). It

recognizes ~50 kDa FIAF from samples of

mouse and rat origins.

Other Names

 ${\sf ANGPTL4}$, ${\sf ANGPTL-4}$, ${\sf UNQ171/PRO197}$, ${\sf FIAF}$, ${\sf ARP4}$, ${\sf ANGPTL2}$, ${\sf Fasting-Induced\ Adipose\ Factor}$, ${\sf NL2}$, ${\sf pp1158}$, ${\sf PSEC0166}$, ${\sf HFARP}$

Target/Specificity

FIAF

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

 $100~\mu g$ (0.5 mg/ml) affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

FIAF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



FIAF Antibody - Protein Information

Name Angptl4

Synonyms Farp, Fiaf {ECO:0000303|PubMed:10862772}

Function

Mediates inactivation of the lipoprotein lipase LPL, and thereby plays a role in the regulation of triglyceride clearance from the blood serum and in lipid metabolism (PubMed:15837923, PubMed:17609370, PubMed:29899519). May also play a role in regulating glucose homeostasis and insulin sensitivity (PubMed:15837923, PubMed:29899519). Inhibits proliferation, migration, and tubule formation of endothelial cells and reduces vascular leakage (PubMed:14583458, PubMed: 17130448, PubMed:21832056). Upon heterologous expression, inhibits the adhesion of endothelial cell to the extracellular matrix (ECM), and inhibits the reorganization of the actin cytoskeleton, formation of actin stress fibers and focal adhesions in endothelial cells that have adhered to ANGPTL4-containing ECM (in vitro) (By similarity). Depending on context, may modulate tumor-related angiogenesis (Probable).

Cellular Location

Secreted. Secreted, extracellular space, extracellular matrix {ECO:0000250|UniProtKB:Q9BY76}. Note=The unprocessed form interacts with the extracellular matrix. This may constitute a dynamic reservoir, a regulatory mechanism of the bioavailability of ANGPTL4. {ECO:0000250|UniProtKB:Q9BY76}

Tissue Location

Detected in liver and kidney (PubMed:10698685, PubMed:17609370). Predominantly expressed in adipose tissue and is strongly up-regulated by fasting in white adipose tissue and liver


FIAF Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FIAF Antibody - Images





Western blot analysis of FIAF expression in rat adipose tissue extract.

FIAF Antibody - Background

Adipose tissue is an energy reserve in animals. Adipose cells produce and secrete numerous physiologically important proteins that help regulate whole body metabolism, such as adiponectin leptin. Fasting causes significant changes in nutrient metabolism and many of these changes are controlled by transcription factors that regulate rate-limiting enzymes. An important transcription factor that mediates metabolic processes induced by fasting is peroxisome proliferator-activated receptor alpha (PPAR alpha). PPAR alpha has recently been reported to target a novel gene encoding the secreted protein FIAF (fasting-induced adipose factor). FIAF is expressed predominantly in adipose tissue and is strongly upregulated in white adipose tissue and the liver under fasting conditions. Initial studies s µggest that FIAF represents a novel endocrine signal helping to regulate metabolism, especially under fasting conditions.