

cAMP Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10417**Specification**

cAMP Antibody - Product Information

Application	E
Reactivity	All Species
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

cAMP Antibody - Additional Information

Application & Usage	1:50 dilutions for nonradioactive EIA assays (10 µl/assay in an 70 µl assay reaction). The material is sufficient for performing 500 nonradioactive EIA assays or 6,000 radioactive assays. Detect 1-100 fmol/assay in a 70 µl assay reaction using nonradioactive method. Refer to the cAMP Assay Kit (K371-100) for detailed assay protocol.
---------------------	---

Other Names
cAMP antibody, Anti- cAMP

Target/Specificity
cAMP

Antibody Form
Liquid

Appearance
Colorless liquid

Formulation
100 µl immunoaffinity purified rabbit IgG in phosphate-buffered saline (PBS) containing 50% glycerol, 1% BSA, and 0.02% sodium azide.

Handling
The antibody solution should be gently mixed before use.

Reconstitution & Storage
-20 °C

Background Descriptions

Precautions
cAMP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

cAMP Antibody - Protein Information

cAMP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

cAMP Antibody - Images

cAMP Antibody - Background

Affinity purified rabbit IgG to 2'O-succinyl adenosine-3',5' cyclic monophosphate (cyclic AMP, cAMP).