

His-Tag Antibody (Clone BV-G020)
Mouse Monoclonal Antibody
Catalog # ABV10489**Specification**

His-Tag Antibody (Clone BV-G020) - Product Information

Application	WB, ICC, IP
Reactivity	All Species
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG2b

His-Tag Antibody (Clone BV-G020) - Additional Information

Application & Usage	Western blotting (1-2 µg/ml), immunoprecipitation (10-20 µg/ml), and immunocytochemistry (10-20 µg/ml). However, the optimal conditions should be determined individually.
---------------------	--

Other Names
Anti His-tag**Target/Specificity**
His-Tag**Antibody Form**
Liquid**Appearance**
Colorless liquid**Formulation**
100 µg (0.5 mg/ml) mouse anti-His-Tag monoclonal antibody in phosphate buffered saline (PBS, pH 7.2), containing 50% glycerol, 1% BSA, 0.02% thimerosal.**Handling**
The antibody solution should be gently mixed before use.**Reconstitution & Storage**
-20 °C**Background Descriptions****Precautions**
His-Tag Antibody (Clone BV-G020) is for research use only and not for use in diagnostic or therapeutic procedures.**His-Tag Antibody (Clone BV-G020) - Protein Information**

His-Tag Antibody (Clone BV-G020) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

His-Tag Antibody (Clone BV-G020) - Images

His-Tag Antibody (Clone BV-G020) - Background

It is expected that the His-Tag monoclonal antibody will react with proteins tagged with His from all species.