10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PIM1 Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10607

Specification

PIM1 Antibody - Product Information

Application	**WB**
Primary Accession	P26794
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**35631**

PIM1 Antibody - Additional Information

Gene ID 24649

Application & Usage	**Western blot analysis (0.5-4 µg/ml). However, the optimal conditions should be determined individually. The antibody recognizes ~33 kDa of PIM1 isoform from samples of human, mouse and rat origins. Reactivity to other species has not been determined.**

Other Names
PIM, PIM1, PIM-1, Proviral Integration Site 1

Target/Specificity
PIM1

Antibody Form
Liquid

Appearance
Colorless liquid

Formulation
100 µg (0.5 mg/ml) affinity purified rabbit anti-PIM1 polyclonal antibody in phosphate (PBS, pH 7.2) containing 30% glycerol, 0.5 % BSA and 0.01% thimerosal.

Handling
The antibody solution should be gently mixed before use.

Reconstitution & Storage
-20 °C

Background Descriptions

Precautions
PIM1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PIM1 Antibody - Protein Information

Name Pim1

Synonyms Pim-1

Function
Proto-oncogene with serine/threonine kinase activity involved in cell survival and cell proliferation and thus providing a selective advantage in tumorigenesis. Exerts its oncogenic activity through: the regulation of MYC transcriptional activity, the regulation of cell cycle progression and by phosphorylation and inhibition of proapoptotic proteins (BAD, MAP3K5, FOXO3) (By similarity). Phosphorylation of MYC leads to an increase of MYC protein stability and thereby an increase of transcriptional activity. The stabilization of MYC exerted by PIM1 might explain partly the strong synergism between these two oncogenes in tumorigenesis. Mediates survival signaling through phosphorylation of BAD, which induces release of the anti-apoptotic protein Bcl- X(L)/BCL2L1 (By similarity). Phosphorylation of MAP3K5, another proapoptotic protein, by PIM1, significantly decreases MAP3K5 kinase activity and inhibits MAP3K5-mediated phosphorylation of JNK and JNK/p38MAPK subsequently reducing caspase-3 activation and cell apoptosis. Stimulates cell cycle progression at the G1-S and G2-M transitions by phosphorylation of CDC25A and CDC25C. Phosphorylation of CDKN1A, a regulator of cell cycle progression at G1, results in the relocation of CDKN1A to the cytoplasm and enhanced CDKN1A protein stability. Promotes cell cycle progression and tumorigenesis by down- regulating expression of a regulator of cell cycle progression, CDKN1B, at both transcriptional and post-translational levels. Phosphorylation of CDKN1B, induces 14-3-3 proteins binding, nuclear export and proteasome-dependent degradation. May affect the structure or silencing of chromatin by phosphorylating HP1 gamma/CBX3 (By similarity). Acts also as a regulator of homing and migration of bone marrow cells involving functional interaction with the CXCL12-CXCR4 signaling axis (By similarity). Acts as a positive regulator of mTORC1 signaling by mediating phosphorylation and inhibition of DEPDC5 component of the GATOR1 complex. Also phosphorylates and activates the ATP-binding cassette transporter ABCG2, allowing resistance to drugs through their excretion from cells (By similarity). Promotes brown adipocyte differentiation (By similarity).

Cellular Location
Cytoplasm. Nucleus. Cell membrane

PIM1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PIM1 Antibody - Images

PIM1 Antibody - Background

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

The proto-oncogene Pim1 belongs to a family of serine/threonine protein kinases that are highly conserved thro µgh evolution in multicellular organisms. Pim1 is involved in the control of cytokine-mediated cell proliferation, differentiation and survival of lymphoid and myeloid cells as well as others. Expression of Pim1 can be stimulated by a variety of growth factors and is regulated at four different levels: transcriptional, post-transcriptional, translational and post-translational. Recently, Pim1 has been shown to enhance the activities of p100, c-Myb and Cdc 25a and in part this might explain reported effects of Pim1 on mitogenesis.