

BLAME Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10692**Specification**

BLAME Antibody - Product Information

Application	WB
Primary Accession	D3ZS24
Other Accession	NP_001099443
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

BLAME Antibody - Additional Information

Application & Usage	Western blotting (0.5-4 µg/ml). However, the optimal conditions should be determined individually. The antibody recognizes ~35 kDa BLAME in Western blot analysis. A 28 kDa cleaved fragment and a 70 kDa glycosylated BLAME can also be detected. Reactivity to other species has not been tested.
---------------------	--

Other Names

B Lymphocyte Activator Macrophage Expressed, Slam family member 8, SLAM-8, SLAMF8

Target/Specificity

BLAME

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions**Precautions**

BLAME Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BLAME Antibody - Protein Information

BLAME Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BLAME Antibody - Images

BLAME Antibody - Background

BLAME (B Lymphocyte Activator Macrophage Expressed) also called SLAMF8. BLAME is a type I transmembrane protein that belongs to the CD2 family of Ig superfamily receptors. BLAME is mostly expressed on macrophages and dendritic cells.