10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

S-Tag Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10758

Specification

S-Tag Antibody - Product Information

Application	**WB**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**

S-Tag Antibody - Additional Information

Application & Usage	**The antibody can be used for Western blot analysis (0.5-4 µg/ml). However, the optimal conditions should be determined individually.**

Other Names
S-tag, anti-S-tag, S-tag antibody

Target/Specificity
S-Tag

Antibody Form
Liquid

Appearance
Colorless liquid

Formulation
100 µg (0.5 mg/ml) affinity purified rabbit anti-S-tag polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, 5mM EDTA and 0.01% thimerosal.

Handling
The antibody solution should be gently mixed before use.

Reconstitution & Storage
-20 °C

Background Descriptions

Precautions
S-Tag Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

S-Tag Antibody - Protein Information


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

S-Tag Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

S-Tag Antibody - Images



Western blot analysis using E.coli lysate expressing S-Tag .

S-Tag Antibody - Background

S-tag is an epitope tag composed of a 15 residue peptide derived from pancreatic ribonuclease A