

GDNF Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10827**Specification**

GDNF Antibody - Product Information

Application	WB, IHC, IP
Primary Accession	P48540
Other Accession	BAA08660
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	23662

GDNF Antibody - Additional Information**Gene ID** 14573

Application & Usage	Western blotting (0.5-4 µg/ml). Based on researcher's feedback, the antibody also works in immunoprecipitation (10-20 µg/ml), and Immunohistochemistry (10-20 µg/ml). However, the optimal concentrations should be determined individually. The antibody recognizes GDNF of human, mouse, and rat origins. Jurkat cell lysate can be used as a positive control.
---------------------	--

Other Names

neurotrophic factor GDNF, GDNF, Glial cell line-derived neurotrophic factor, Astrocyte-derived trophic factor; ATF

Target/Specificity

GDNF

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.2 mg/ml) protein A affinity purified rabbit polyclonal antibody in phosphate-buffered saline (PBS) containing 30% glycerol, 0.5% BSA, and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

GNDF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

GNDF Antibody - Protein Information

Name Gdnf

Function

Neurotrophic factor that enhances survival and morphological differentiation of dopaminergic neurons and increases their high- affinity dopamine uptake.

Cellular Location

Secreted {ECO:0000250|UniProtKB:P39905}.

Tissue Location

Expressed in both the central nervous system (CNS) and in non-CNS tissues. Expressed in a highly dynamic pattern in the anterior neuroectoderm during the early stages of neurogenesis between 7.5 dpc and 10.5 dpc. Beginning at 10.5 dpc, expression begins in mesenchymal tissues of several organs including the digestive tract, kidney, testis, frontonasal mass, tooth primordium, tongue, mandible, whisker follicles, ear, eye, limb bud and in distinct regions of the brain. Also expressed in the heart, ileum, liver and muscle

GNDF Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GNDF Antibody - Images

GNDF Antibody - Background

Glial cell line-derived neurotrophic factor (GNDF) has been identified as a potent neurotrophic factor that enhances survival of midbrain dopaminergic neurons. GNDF is a glycosylated, disulfide-bonded homodimer. It promotes the survival and morphological differentiation of dopaminergic neurons and increases their high-affinity dopamine uptake. On the basis of these findings, it has been suggested that GNDF may play a role in the treatment of Parkinson's disease, which is marked by progressive degeneration of midbrain dopaminergic neurons.