

FABP5 Antibody
Rabbit Polyclonal Antibody
Catalog # ABV10982**Specification**

FABP5 Antibody - Product Information

Application	WB
Primary Accession	Q01469
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	15164

FABP5 Antibody - Additional Information**Gene ID** 2171

Positive Control	Recombinant human FABP-5
Application & Usage	Western Blot analysis (0.5-4 µg/ml). However, the optimal conditions should be determined individually. Recombinant human FABP-5 can be used as a positive control.

Other Names

Epidermal-type fatty acid-binding protein, E-FABP, Fatty acid-binding protein 5, Psoriasis-associated fatty acid-binding protein homolog

Target/Specificity

FABP5

Antibody Form

Liquid

Appearance

Colorless liquid

Formulation

100 µg (0.5 mg/ml) affinity purified rabbit anti-human FABP5 polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 30% glycerol, 0.5% BSA, 5mM EDTA and 0.01% thimerosal.

Handling

The antibody solution should be gently mixed before use.

Reconstitution & Storage

-20 °C

Background Descriptions

Precautions

FABP5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

FABP5 Antibody - Protein Information

Name FABP5 ([HGNC:3560](#))

Function

Intracellular carrier for long-chain fatty acids and related active lipids, such as endocannabinoids, that regulate the metabolism and actions of the ligands they bind. In addition to the cytosolic transport, selectively delivers specific fatty acids from the cytosol to the nucleus, wherein they activate nuclear receptors (PubMed:22170058, PubMed:21395585). Delivers retinoic acid to the nuclear receptor peroxisome proliferator-activated receptor delta; which promotes proliferation and survival. May also serve as a synaptic carrier of endocannabinoid at central synapses and thus controls retrograde endocannabinoid signaling. Modulates inflammation by regulating PTGES induction via NF-kappa-B activation, and prostaglandin E2 (PGE2) biosynthesis during inflammation (By similarity). May be involved in keratinocyte differentiation (PubMed:8092987).

Cellular Location

Cytoplasm. Nucleus Synapse {ECO:0000250|UniProtKB:Q05816}. Postsynaptic density {ECO:0000250|UniProtKB:Q05816}. Secreted {ECO:0000250|UniProtKB:Q05816}. Note=Localizes primarily to the cytoplasm. Upon certain ligand binding, a conformation change exposes a nuclear localization motif and the protein is transported into nucleus (PubMed:24692551). Secreted by astrocytes, but not by neurons (By similarity). {ECO:0000250|UniProtKB:Q05816, ECO:0000269|PubMed:24692551}

Tissue Location

Keratinocytes; highly expressed in psoriatic skin (PubMed:8092987). Expressed in brain gray matter (PubMed:21395585)

FABP5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FABP5 Antibody - Images**FABP5 Antibody - Background**

FABPs have been reported to be involved in the uptake and metabolism of fatty acids, maintenance of cellular membrane fatty acids levels, intracellular trafficking, modulation of specific enzymes of lipid metabolic pathways, as well as in the modulation of cell growth and differentiation.