

## LIGHT Antibody

Rabbit Polyclonal Antibody Catalog # ABV11019

## Specification

# LIGHT Antibody - Product Information

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW

O43557 AAC39563 Human Rabbit Polyclonal Rabbit IgG 26350

WB

# LIGHT Antibody - Additional Information

Gene ID 8740

Application & Usage

Western blotting (0.5-1  $\mu$ g/ml). However, the optimal conditions should be determined individually. The antibody detects human LIGHT. Reactivity to other species has not been tested.

Other Names Tumor necrosis factor ligand, TNF14, TNF 14, TNF-14

Target/Specificity LIGHT

Antibody Form Liquid

Appearance Colorless liquid

**Formulation** 100  $\mu$ g (0.5 mg/ml) antigen affinity purified rabbit polyclonal antibody in phosphate buffered saline (PBS), pH 7.2, containing 50% glycerol, 1% BSA, 0.02% thimerosal.

Handling The antibody solution should be gently mixed before use.

Reconstitution & Storage -20 °C

**Background Descriptions** 

Precautions



LIGHT Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# LIGHT Antibody - Protein Information

Name TNFSF14

Synonyms HVEML, LIGHT

#### Function

Cytokine that binds to TNFRSF3/LTBR. Binding to the decoy receptor TNFRSF6B modulates its effects. Acts as a ligand for TNFRSF14/HVEM (PubMed:<a href="http://www.uniprot.org/citations/9462508" target="\_blank">9462508</a>, PubMed:<a href="http://www.uniprot.org/citations/10754304" target="\_blank">10754304</a>). Upon binding to TNFRSF14/HVEM, delivers costimulatory signals to T cells, leading to T cell proliferation and IFNG production (PubMed:<a href="http://www.uniprot.org/citations/10754304" target="\_blank">10754304" target="\_blank">10754304</a>).

#### **Cellular Location**

[Tumor necrosis factor ligand superfamily member 14, membrane form]: Cell membrane; Single-pass type II membrane protein [Isoform 2]: Cytoplasm.

#### **Tissue Location**

Predominantly expressed in the spleen but also found in the brain. Weakly expressed in peripheral lymphoid tissues and in heart, placenta, liver, lung, appendix, and kidney, and no expression seen in fetal tissues, endocrine glands, or nonhematopoietic tumor lines.

## LIGHT Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# LIGHT Antibody - Images

## LIGHT Antibody - Background

LIGHT is a type II membrane protein belonging to the TNF family of membrane-anchored ligands. It is expressed in splenocytes, activated PBL, CD8+ tumor infiltrating lymphocytes, granulocytes and monocytes. LIGHT has the ability to activate NFkB, stimulate proliferation of lymphocytes, and induce apoptosis in certain human tumor cells. LIGHT binds to three receptors; the herpes virus entry mediator type A receptor (HEVM), LTbR and a decoy receptor (DcR3).