

**H3K4me3 polyclonal antibody**  
**Rabbit Polyclonal Antibody**  
**Catalog # ABV11399****Specification**

---

**H3K4me3 polyclonal antibody - Product Information**

Application	<b>CHIP, DB, WB</b>
Primary Accession	<a href="#">P68431</a>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>15404</b>

**H3K4me3 polyclonal antibody - Additional Information****Gene ID** 8350;8351;8352;8353;8354;8355;8356;8357;8358;8968

Positive Control	<b>Western blot: HeLa cells, IF: HeLa cells, ELISA: Antigen, ChIP: HeLa cells, Dot blot: Histone Peptides.</b>
Application & Usage	<b>WB: 1:500, Dot Blot: 1:2,000, ChIP: 1 µl/ChIP, IF: 1:500.</b>

**Other Names**  
Histone 3**Target/Specificity**  
H3K4me3**Antibody Form**  
Liquid**Appearance**  
Colorless liquid**Formulation**  
In PBS with 0.05% sodium azide and 0.05% ProClin 300.**Handling**  
The antibody solution should be gently mixed before use.**Reconstitution & Storage**  
-20 °C**Background Descriptions****Precautions**  
H3K4me3 polyclonal antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## H3K4me3 polyclonal antibody - Protein Information

**Name** H3C1 ([HGNC:4766](#))

**Synonyms** H3FA, HIST1H3A

### Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

### Cellular Location


Nucleus. Chromosome.

## H3K4me3 polyclonal antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## H3K4me3 polyclonal antibody - Images



ChIP assays were performed using HeLa cells and the antibody and optimized PCR primer sets for qPCR. A titration of the antibody consisting of 1, 2, 5, and 10 µl per ChIP experiment was analysed. IgG (2 µg/IP) was used as negative control. The Fig shows the recovery, expressed as a % of input (the relative amount of IP DNA compared to input DNA after qPCR analysis).



A Dot Blot analysis was performed to test the cross reactivity of the antibody with peptides containing other modifications of histone H4 and H3. 100 to 0.2 pmol of the peptide containing the respective histone modification were spotted on a membrane. The Fig shows a high specificity of the antibody for the modification of interest.



HeLa cells (15 µg) were analysed by WB blot using the antibody.

### H3K4me3 polyclonal antibody - Background

Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histone tails undergo numerous post-translational modifications, which either directly or indirectly alter chromatin structure to facilitate transcriptional activation or repression or other nuclear processes. In addition to the genetic code, combinations of the different histone modifications reveal the so-called "histone code". Histone methylation and demethylation is dynamically regulated by respectively histone methyl transferases and histone demethylases. Trimethylation of histone H3K4 is associated with active promoters.