

TIP60 Antibody
Purified Rabbit Polyclonal Antibody
Catalog # ABV11625**Specification**

TIP60 Antibody - Product Information

Application	WB, IHC, IF
Primary Accession	O92993
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	58582

TIP60 Antibody - Additional Information**Gene ID** 10524**Other Names**

KAT5 Antibody: lysine acetyltransferase 5, K-acetyltransferase 5, lysine acetyltransferase 5, HIV-1 Tat interactive protein 60kDa, HTATIP, HTATIP1, TIP, cPLA2 interacting protein, cPLA2, ESA1, PLIP

Target/Specificity

TIP60

Formulation

1 mg/ml in PBS containing 0.02% sodium azide

Handling

The antibody solution should be gently mixed before use.

Background Descriptions**Precautions**

TIP60 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TIP60 Antibody - Protein Information**Name** KAT5 {ECO:0000303|PubMed:32817552, ECO:0000312|HGNC:HGNC:5275}**Function**

Catalytic subunit of the NuA4 histone acetyltransferase complex, a multiprotein complex involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H2A and H4 (PubMed: [12776177](http://www.uniprot.org/citations/12776177), PubMed: [15042092](http://www.uniprot.org/citations/15042092), PubMed: [15121871](http://www.uniprot.org/citations/15121871), PubMed: [15310756](http://www.uniprot.org/citations/15310756))

target="_blank">15310756, PubMed:14966270, PubMed:16387653, PubMed:19909775, PubMed:25865756, PubMed:27153538, PubMed:29335245, PubMed:29174981, PubMed:33076429, PubMed:32822602). Histone acetylation alters nucleosome-DNA interactions and promotes interaction of the modified histones with other proteins which positively regulate transcription (PubMed:12776177, PubMed:15042092, PubMed:15121871, PubMed:15310756, PubMed:14966270). The NuA4 histone acetyltransferase complex is required for the activation of transcriptional programs associated with proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair (PubMed:17709392, PubMed:19783983, PubMed:32832608). The NuA4 complex plays a direct role in repair of DNA double-strand breaks (DSBs) by promoting homologous recombination (HR): the complex inhibits TP53BP1 binding to chromatin via MBTD1, which recognizes and binds histone H4 trimethylated at 'Lys-20' (H4K20me), and KAT5 that catalyzes acetylation of 'Lys-15' of histone H2A (H2AK15ac), thereby blocking the ubiquitination mark required for TP53BP1 localization at DNA breaks (PubMed:27153538, PubMed:32832608). Also involved in DSB repair by mediating acetylation of 'Lys-5' of histone H2AX (H2AXK5ac), promoting NBN/NBS1 assembly at the sites of DNA damage (PubMed:17709392, PubMed:26438602). The NuA4 complex plays a key role in hematopoietic stem cell maintenance and is required to maintain acetylated H2A.Z/H2AZ1 at MYC target genes (By similarity). The NuA4 complex is also required for spermatid development by promoting acetylation of histones: histone hyperacetylation is required for histone replacement during the transition from round to elongating spermatids (By similarity). Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AZ1 from the nucleosome (PubMed:24463511). Also acetylates non-histone proteins, such as BMAL1, ATM, AURKB, CHKA, CGAS, ERCC4/XPF, LPIN1, NDC80/HEC1, NR1D2, RAN, SOX4, FDXP3, SQSTM1, ULK1 and RUBCNL/Pacer (PubMed:16141325, PubMed:17360565, PubMed:17996965, PubMed:24835996, PubMed:26829474, PubMed:29040603, PubMed:30409912, PubMed:30704899, PubMed:31857589, PubMed:32034146, PubMed:32817552, PubMed:34077757). Directly acetylates and activates ATM (PubMed:16141325). Promotes nucleotide excision repair (NER) by mediating acetylation of ERCC4/XPF, thereby promoting formation of the ERCC4-ERCC1 complex (PubMed:<a

[32034146](http://www.uniprot.org/citations/32034146)). Relieves NR1D2-mediated inhibition of APOC3 expression by acetylating NR1D2 (PubMed:[17996965](http://www.uniprot.org/citations/17996965)). Acts as a regulator of regulatory T-cells (Treg) by catalyzing FOXP3 acetylation, thereby promoting FOXP3 transcriptional repressor activity (PubMed:[17360565](http://www.uniprot.org/citations/17360565)), PubMed:[24835996](http://www.uniprot.org/citations/24835996)). Involved in skeletal myoblast differentiation by mediating acetylation of SOX4 (PubMed:[26291311](http://www.uniprot.org/citations/26291311)). Catalyzes acetylation of APBB1/FE65, increasing its transcription activator activity (PubMed:[33938178](http://www.uniprot.org/citations/33938178)). Promotes transcription elongation during the activation phase of the circadian cycle by catalyzing acetylation of BMAL1, promoting elongation of circadian transcripts (By similarity). Together with GSK3 (GSK3A or GSK3B), acts as a regulator of autophagy: phosphorylated at Ser-86 by GSK3 under starvation conditions, leading to activate acetyltransferase activity and promote acetylation of key autophagy regulators, such as ULK1 and RUBCNL/Pacer (PubMed:[30704899](http://www.uniprot.org/citations/30704899)). Acts as a regulator of the cGAS-STING innate antiviral response by catalyzing acetylation the N-terminus of CGAS, thereby promoting CGAS DNA-binding and activation (PubMed:[32817552](http://www.uniprot.org/citations/32817552)). Also regulates lipid metabolism by mediating acetylation of CHKA or LPIN1 (PubMed:[34077757](http://www.uniprot.org/citations/34077757)). Promotes lipolysis of lipid droplets following glucose deprivation by mediating acetylation of isoform 1 of CHKA, thereby promoting monomerization of CHKA and its conversion into a tyrosine-protein kinase (PubMed:[34077757](http://www.uniprot.org/citations/34077757)). Acts as a regulator of fatty-acid-induced triacylglycerol synthesis by catalyzing acetylation of LPIN1, thereby promoting the synthesis of diacylglycerol (PubMed:[29765047](http://www.uniprot.org/citations/29765047)). In addition to protein acetyltransferase, can use different acyl-CoA substrates, such as (2E)-butenoyl-CoA (crotonyl-CoA) and 2-hydroxyisobutanoyl-CoA (2-hydroxyisobutyryl-CoA), and is able to mediate protein crotonylation and 2-hydroxyisobutyrylation, respectively (PubMed:[29192674](http://www.uniprot.org/citations/29192674)), PubMed:[34608293](http://www.uniprot.org/citations/34608293)). Acts as a key regulator of chromosome segregation and kinetochore-microtubule attachment during mitosis by mediating acetylation or crotonylation of target proteins (PubMed:[26829474](http://www.uniprot.org/citations/26829474)), PubMed:[29040603](http://www.uniprot.org/citations/29040603), PubMed:[30409912](http://www.uniprot.org/citations/30409912), PubMed:[34608293](http://www.uniprot.org/citations/34608293)). Catalyzes acetylation of AURKB at kinetochores, increasing AURKB activity and promoting accurate chromosome segregation in mitosis (PubMed:[26829474](http://www.uniprot.org/citations/26829474)). Acetylates RAN during mitosis, promoting microtubule assembly at mitotic chromosomes (PubMed:[29040603](http://www.uniprot.org/citations/29040603)). Acetylates NDC80/HEC1 during mitosis, promoting robust kinetochore- microtubule attachment (PubMed:[30409912](http://www.uniprot.org/citations/30409912)). Catalyzes crotonylation of MAPRE1/EB1, thereby ensuring accurate spindle positioning in mitosis (PubMed:[34608293](http://www.uniprot.org/citations/34608293)).

Cellular Location

Nucleus. Chromosome. Cytoplasm Chromosome, centromere, kinetochore Cytoplasm, cytoskeleton, spindle pole Nucleus, nucleolus. Cytoplasm, perinuclear region. Note=Upon stimulation with EDN1, it is exported from the nucleus to the perinuclear region and UV irradiation induces translocation into punctuate subnuclear structures named nuclear bodies (PubMed:11262386). Transiently localizes to kinetochores in early mitosis (PubMed:26829474). Localizes to spindle poles when chromosomes align during metaphase (PubMed:34608293). Localizes in the cytoplasm and nucleus of round spermatids (By similarity).

{ECO:0000250|UniProtKB:Q8CHK4, ECO:0000269|PubMed:11262386,
ECO:0000269|PubMed:26829474, ECO:0000269|PubMed:34608293}

TIP60 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TIP60 Antibody - Images

TIP60 Antibody - Background

TIP60 belongs to the MYST family of histone acetyl transferases (HATs) and was originally isolated as an HIV-1 TAT interactive protein. HATs play important roles in regulating chromatin remodeling, transcription and other nuclear processes by acetylating histone and non-histone proteins. TIP60 has been shown to play a role in DNA repair and apoptosis and is thought to play an important role in signal transduction and autophagy. In case of HIV-1 infection, interaction with the viral Tat protein leads to TIP60 polyubiquitination and targets it to degradation. Also, TIP60 contains a chromodomain. TIP60 is a catalytic subunit of the NuA4 histone acetyltransferase complex. The NuA4 HAT complex plays a role in transcriptional activation of select genes mainly by acetylation of nucleosomal histone H4 and H2A, which influence nucleosome-DNA interaction and promotes interaction of the modified histones with other proteins that could regulate transcription positively. This complex also involves in the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, DNA repair and apoptosis.